

EENS 2120

Petrology

Prof. Stephen A. Nelson

Types of Metamorphism

This document last updated on 12-Apr-2018

Metamorphism is defined as follows:

The mineralogical and structural adjustment of solid rocks to physical and chemical conditions that have been imposed at depths below the near surface zones of weathering and diagenesis and which differ from conditions under which the rocks in question originated.

The word "*Metamorphism*" comes from the Greek: meta = after, morph = form, so metamorphism means the after form. In geology this refers to the changes in mineral assemblage and texture that result from subjecting a rock to conditions such pressures, temperatures, and chemical environments different from those under which the rock originally formed.


- Note that *Diagenesis* is also a change in form that occurs in sedimentary rocks. In geology, however, we restrict diagenetic processes to those which occur at temperatures below 200°C and pressures below about 300 MPa (MPa stands for Mega Pascals), this is equivalent to about 3 kilobars of pressure (1kb = 100 MPa).
- Metamorphism, therefore occurs at temperatures and pressures higher than 200°C and 300 MPa. Rocks can be subjected to these higher temperatures and pressures as they are buried deeper in the Earth. Such burial usually takes place as a result of tectonic processes such as continental collisions or subduction.
- The upper limit of metamorphism occurs at the pressure and temperature where melting of the rock in question begins. Once melting begins, the process changes to an igneous process rather than a metamorphic process.

Grade of Metamorphism

As the temperature and/or pressure increases on a body of rock we say the rock undergoes *prograde metamorphism* or that the grade of metamorphism increases. *Metamorphic grade* is a general term for describing the relative temperature and pressure conditions under which metamorphic rocks form.

- Low-grade metamorphism takes place

at temperatures between about 200 to 320°C, and relatively low pressure. Low grade metamorphic rocks are generally characterized by an abundance of hydrous minerals. With increasing grade of metamorphism, the hydrous minerals begin to react with other minerals and/or break down to less hydrous minerals.




- High-grade metamorphism takes place at temperatures greater than 320°C and relatively high pressure. As grade of metamorphism increases, hydrous minerals become less hydrous, by losing H₂O, and non-hydrous minerals become more common.

Types of Metamorphism

Contact Metamorphism

Contact metamorphism occurs adjacent to igneous intrusions and results from high temperatures associated with the igneous intrusion.

Since only a small area surrounding the intrusion is heated by the magma, metamorphism is restricted to the zone surrounding the intrusion, called a **metamorphic** or **contact aureole**. Outside of the contact aureole, the rocks are not affected by the intrusive event. The grade of metamorphism increases in all directions toward the intrusion. Because the temperature contrast between the surrounding rock and the intruded magma is larger at shallow levels in the crust where pressure is low, contact metamorphism is often referred to as high temperature, low pressure metamorphism. The rock produced is often a fine-grained rock that shows no foliation, called a **hornfels**.



Regional Metamorphism

Regional metamorphism occurs over large areas and generally does not show any relationship to igneous bodies. Most regional metamorphism is accompanied by deformation under non-hydrostatic or differential stress conditions. Thus, regional metamorphism usually results in forming metamorphic rocks that are strongly foliated, such as slates, schists, and gniesses. The differential stress usually results from tectonic forces that produce compressional stresses in the rocks, such as when two continental masses collide. Thus, regionally metamorphosed rocks occur in the cores of fold/thrust mountain belts or in eroded mountain ranges. Compressive stresses result in folding of rock and thickening of the crust, which tends to push rocks to deeper levels where they are subjected to higher temperatures and pressures.

Cataclastic Metamorphism

Cataclastic metamorphism occurs as a result of mechanical deformation, like when two bodies of rock slide past one another along a fault zone. Heat is generated by the friction of sliding along such a shear zone, and the rocks tend to be mechanically deformed, being crushed and pulverized, due to the shearing. Cataclastic metamorphism is not very common and is restricted to a narrow zone along which the shearing occurred.

Hydrothermal Metamorphism

Rocks that are altered at high temperatures and moderate pressures by hydrothermal fluids are hydrothermally metamorphosed. This is common in basaltic rocks that generally lack hydrous minerals. The hydrothermal metamorphism results in alteration to such Mg-Fe rich hydrous minerals as talc, chlorite, serpentine, actinolite, tremolite, zeolites, and clay minerals. Rich ore deposits are often formed as a result of hydrothermal metamorphism.

Burial Metamorphism

When sedimentary rocks are buried to depths of several kilometers, temperatures greater than 300°C may develop in the absence of differential stress. New minerals grow, but the rock does not appear to be metamorphosed. The main minerals produced are often the Zeolites. Burial metamorphism overlaps, to some extent, with diagenesis, and grades into regional metamorphism as temperature and pressure increase.

Shock Metamorphism (Impact Metamorphism)

When an extraterrestrial body, such as a meteorite or comet impacts with the Earth or if there is a very large volcanic explosion, ultrahigh pressures can be generated in the impacted rock. These ultrahigh pressures can produce minerals that are only stable at very high pressure, such as the SiO₂ polymorphs coesite and stishovite. In addition they can produce textures known as shock lamellae in mineral grains, and such textures as shatter cones in the impacted rock.

Classification of Metamorphic Rocks

Classification of metamorphic rocks is based on mineral assemblage, texture, protolith, and bulk chemical composition of the rock. Each of these will be discussed in turn, then we will summarize how metamorphic rocks are classified.

Texture

In metamorphic rocks individual minerals may or may not be bounded by crystal faces. Those that are bounded by their own crystal faces are termed *idioblastic*. Those that show none of their own crystal faces are termed *xenoblastic*. From examination of metamorphic rocks, it has been found that metamorphic minerals can be listed in a generalized sequence, known as the *crystalloblastic series*, listing minerals in order of their tendency to be idioblastic. In the series, each mineral tends to develop idioblastic surfaces against any mineral that occurs lower in the series. This series is listed below:

- rutile, sphene, magnetite
- tourmaline kyanite, staurolite, garnet, andalusite
- epidote, zoisite, lawsonite, forsterite
- pyroxenes, amphiboles, wollastonite
- micas, chlorites, talc, stilpnomelane, prehnite
- dolomite, calcite
- scapolite, cordierite, feldspars
- quartz

This series can, in a rather general way, enable us to determine the origin of a given rock. For example a rock that shows euhedral plagioclase crystals in contact with anhedral amphibole, likely had an igneous protolith, since a metamorphic rock with the same minerals would be expected to show euhedral amphibole in contact with anhedral plagioclase.

Another aspect of the crystalloblastic series is that minerals high on the list tend to form **porphyroblasts** (the metamorphic equivalent of phenocrysts), although K-feldspar (a mineral that occurs lower in the list) may also form porphyroblasts. Porphyroblasts are often riddled with inclusions of other minerals that were enveloped during growth of the porphyroblast. These are said to have a **poikioblastic texture**.

Most metamorphic textures involve foliation. Foliation is generally caused by a preferred orientation of sheet silicates. If a rock has a slaty cleavage as its foliation, it is termed a **slate**, if it has a phyllitic foliation, it is termed a **phyllite**, if it has a shistose foliation, it is termed a **schist**. A rock that shows a banded texture without a distinct foliation is termed a **gneiss**. All of these could be porphyroblastic (i.e. could contain porphyroblasts).

A rock that shows no foliation is called a **hornfels** if the grain size is small, and a **granulite**, if the grain size is large and individual minerals can be easily distinguished with a hand lens.

Protolith

Protolith refers to the original rock, prior to metamorphism. In low grade metamorphic rocks, original textures are often preserved allowing one to determine the likely protolith. As the grade of metamorphism increases, original textures are replaced with metamorphic textures and other clues, such as bulk chemical composition of the rock, are used to determine the protolith.

Bulk Chemical Composition

The mineral assemblage that develops in a metamorphic rock is dependent on

- The pressure and temperature reached during metamorphism
- The composition of any fluid phase present during metamorphism, and
- The bulk chemical composition of the rock.

Just like in igneous rocks, minerals can only form if the necessary chemical constituents are present in the rock (i.e. the concept of silica saturation and alumina saturation applies to metamorphic rocks as well). Based on the mineral assemblage present in the rock one can often estimate the approximate bulk chemical composition of the rock. Some terms that describe this general bulk chemical composition are as follows:

- **Pelitic**. These rocks are derivatives of aluminous sedimentary rocks like shales and mudrocks. Because of their high concentrations of alumina they are recognized by an abundance of aluminous minerals, like clay minerals, micas, kyanite, sillimanite, andalusite, and garnet.
- **Quartzo-Feldspathic**. Rocks that originally contained mostly quartz and feldspar like granitic rocks and arkosic sandstones will also contain an abundance of quartz and feldspar as metamorphic rocks, since these minerals are stable over a wide range of temperature and pressure. Those that exhibit mostly quartz and feldspar with only minor

amounts of aluminous minerals are termed quartzo-feldspathic.

- **Calcareous.** Calcareous rocks are calcium rich. They are usually derivatives of carbonate rocks, although they contain other minerals that result from reaction of the carbonates with associated siliceous detrital minerals that were present in the rock. At low grades of metamorphism calcareous rocks are recognized by their abundance of carbonate minerals like calcite and dolomite. With increasing grade of metamorphism these are replaced by minerals like brucite, phlogopite (Mg-rich biotite), chlorite, and tremolite. At even higher grades anhydrous minerals like diopside, forsterite, wollastonite, grossularite, and calcic plagioclase.
- **Basic.** Just like in igneous rocks, the general term basic refers to low silica content. Basic metamorphic rocks are generally derivatives of basic igneous rocks like basalts and gabbros. They have an abundance of Fe-Mg minerals like biotite, chlorite, and hornblende, as well as calcic minerals like plagioclase and epidote.
- **Magnesian.** Rocks that are rich in Mg with relatively less Fe, are termed magnesian. Such rocks would contain Mg-rich minerals like serpentine, brucite, talc, dolomite, and tremolite. In general, such rocks usually have an ultrabasic protolith, like peridotite, dunite, or pyroxenite.
- **Ferruginous.** Rocks that are rich in Fe with little Mg are termed ferruginous. Such rocks could be derivatives of Fe-rich cherts or ironstones. They are characterized by an abundance of Fe-rich minerals like greenalite (Fe-rich serpentine), minnesotaite (Fe-rich talc), ferroactinolite, ferrocummingtonite, hematite, and magnetite at low grades, and ferrosilite, fayalite, ferrohedenbergite, and almandine garnet at higher grades.
- **Manganiferrous.** Rocks that are characterized by the presence of Mn-rich minerals are termed manganiferrous. They are characterized by such minerals as Stilpnomelane and spessartine.

Classification

Classification of metamorphic rocks depends on what is visible in the rock and its degree of metamorphism. Note that classification is generally loose and practical such that names can be adapted to describe the rock in the most satisfactory way that conveys the important characteristics. Three kinds of criteria are normally employed. These are:

1. Mineralogical - The most distinguishing minerals are used as a prefix to a textural term. Thus, a schist containing biotite, garnet, quartz, and feldspar, would be called a biotite-garnet schist. A gneiss containing hornblende, pyroxene, quartz, and feldspar would be called a hornblende-pyroxene gneiss. A schist containing porphyroblasts of K-feldspar would be called a K-spar porphyroblastic schist.
2. Chemical - If the general chemical composition can be determined from the mineral assemblage, then a chemical name can be employed. For example a schist with a lot of quartz and feldspar and some garnet and muscovite would be called a garnet-muscovite quartzo-feldspathic schist. A schist consisting mostly of talc would be called a talc-magnesian schist.
3. Protolithic - If a rock has undergone only slight metamorphism such that its original texture can still be observed then the rock is given a name based on its original name, with the prefix meta- applied. For example: metabasalt, metagraywacke, meta-andesite,

metagranite.

In addition to these conventions, certain non-foliated rocks with specific chemical compositions and/or mineral assemblages are given specific names. These are as follows:


- ***Amphibolites***: These are medium to coarse grained, dark colored rocks whose principal minerals are hornblende and plagioclase. They result from metamorphism of basic igneous rocks. Foliation is highly variable, but when present the term schist can be appended to the name (i.e. amphibolite schist).
- ***Marbles***: These are rocks composed mostly of calcite, and less commonly of dolomite. They result from metamorphism of limestones and dolostones. Some foliation may be present if the marble contains micas.
- ***Eclogites***: These are medium to coarse grained consisting mostly of garnet and green clinopyroxene called omphacite, that result from high grade metamorphism of basic igneous rocks. Eclogites usually do not show foliation.
- ***Quartzites***: Quartz arenites and chert both are composed mostly of SiO_2 . Since quartz is stable over a wide range of pressures and temperatures, metamorphism of quartz arenites and cherts will result only in the recrystallization of quartz forming a hard rock with interlocking crystals of quartz. Such a rock is called a quartzite.
- ***Serpentinites***: Serpentinites are rocks that consist mostly of serpentine. These form by hydrothermal metamorphism of ultrabasic igneous rocks.
- ***Soapstones***: Soapstones are rocks that contain an abundance of talc, which gives the rock a greasy feel, similar to that of soap. Talc is an Mg-rich mineral, and thus soapstones from ultrabasic igneous protoliths, like peridotites, dunites, and pyroxenites, usually by hydrothermal alteration.
- ***Skarns***: Skarns are rocks that originate from contact metamorphism of limestones or dolostones, and show evidence of having exchanged constituents with the intruding magma. Thus, skarns are generally composed of minerals like calcite and dolomite, from the original carbonate rock, but contain abundant calcium and magnesium silicate minerals like andradite, grossularite, epidote, vesuvianite, diopside, and wollastonite that form by reaction of the original carbonate minerals with silica from the magma. The chemical exchange that takes place is called ***metasomatism***.
- ***Mylonites***: Mylonites are cataclastic metamorphic rocks that are produced along shear zones deep in the crust. They are usually fine-grained, sometimes glassy, that are streaky or layered, with the layers and streaks having been drawn out by ductile shear.

Metamorphic Facies

In general, metamorphic rocks do not drastically change chemical composition during metamorphism, except in the special case where metasomatism is involved (such as in the production of skarns, as discussed above). The changes in mineral assemblages are due to changes in the temperature and pressure conditions of metamorphism. Thus, the mineral assemblages that are observed must be an indication of the temperature and pressure

environment that the rock was subjected to. This pressure and temperature environment is referred to as *Metamorphic Facies*. (This is similar to the concept of sedimentary facies, in that a sedimentary facies is also a set of environmental conditions present during deposition). The sequence of metamorphic facies observed in any metamorphic terrain, depends on the geothermal gradient that was present during metamorphism.

A high geothermal gradient such as the one labeled "A", might be present around an igneous intrusion, and would result in metamorphic rocks belonging to the hornfels facies. Under a normal to high geothermal gradient, such as "B", rocks would progress from zeolite facies to greenschist, amphibolite, and eclogite facies as the grade of metamorphism (or depth of burial) increased. If a low geothermal gradient was present, such the one labeled "C" in the diagram, then rocks would progress from zeolite facies to blueschist facies to eclogite facies.



Thus, if we know the facies of metamorphic rocks in the region, we can determine what the geothermal gradient must have been like at the time the metamorphism occurred. This relationship between geothermal gradient and metamorphism will be the central theme of our discussion of metamorphism.

Examples of questions on this material that could be asked on an exam

1. Briefly define each of the seven types of metamorphism.
 2. Explain how metamorphic rocks are classified.
 3. What is the crystalloblastic series?
 4. Which minerals commonly form porphyroblasts?
 5. Define the following metamorphic rocks: (a) quartzite, (b) marble, (c) eclogite, (d) amphibolite, (e) soapstone (f) serpentinite, (g) mylonite
 6. What is a geothermal gradient?
-

[Return to EENS 2120 Home Page](#)